

POSTER PRESENTATION

Open Access

Antimicrobial potential of *Actinomycetes* by NRPS and PKS-I pathways

Maria Lucila Hernández- Macedo^{1*}, Erick Nunes Barreto¹, Ana Carolina de Souza Cavalcanti², Rafael Salomão da Silva², Erika Cristina Teixeira dos Anjos Brandão², Roberta Pereira Miranda Fernandes³, Viviane Talamini², Leandro Eugenio Cardamone Diniz⁴, Marcelo Ferreira Fernandes²

From 5th Congress of the Brazilian Biotechnology Society (SBBIOTEC)
Florianópolis, Brazil. 10-14 November 2013

Background

Actinomycetes may account for 10 to 30% of the total soil rhizosphere microorganisms. The attention given to the actinomycetes in biotechnological applications is a result of their metabolic versatility that is accompanied by the production of primary and secondary metabolites of economic importance, which are a promising source of products (*e.g.*, antibiotics, enzyme inhibitors, antiparasitic and anticancer agents) [1,2]. Included in this range of compounds are secondary metabolites synthesized by polyketide synthase (PKS) and non-ribosomal peptide synthetase (NRPS) pathways. An effective method for assessing the presence of these biosynthetic pathways is the detection of PKS and NRPS genes by PCR [3,4]. Thus, this study was based on targeted analyses of 31 soil isolate actinomycetes aiming to evaluate their antimicrobial potential through the NRPS and PKS-I pathways.

Methods

The antimicrobial activity was evaluated by the antagonism test against two economically important phytopathogens, the bacterium *Xanthomonas campestris* and the fungus *Thielaviopsis paradoxa*, using the technique of double layer. The *X. campestris* and *T. paradoxa* were propagated at 28°C in YM (yeast malt) pH 6.0 and PDB (potato dextrose broth) respectively. The results were statistically analyzed using the Bonferroni test.

The presence of genes PKS and NRPS was evaluated by PCR, using degenerate primers for highly conserved regions encoding enzymes associated with biosynthesis of polyketides and peptides.

Results and conclusions

According to the experimental results, 52% of the isolates showed antimicrobial activity against at least one of the target bacterial pathogens tested. Among these active isolates, some belong to rare families. Thus, this finding can be a source of novel biomolecules with antimicrobial activity. From those isolates that presented one of the NRPS and PKS-I genes, 75% of them showed antagonistic activity against one of the phytopathogens evaluated. Preliminary data on this screening demonstrate the importance of the biotechnological potential of these actinomycetes due to the antagonistic activity against plant pathogens of economic interest and the possibility of be used as biocontrol, besides offering a strong area for metabolic research [2,5].

Acknowledgements

CNPq and FAPITEC for financial support and fellowship, and Embrapa-Tabuleiros Costeiros, SE, Brazil (Laboratório de Biotecnologia e Biologia Molecular and Laboratório de Microbiologia do Solo for technical support.

Authors' details

¹Programa de Pós-Graduação em Biotecnologia Industrial, Universidade Tiradentes/Instituto de Tecnologia e Pesquisa, Aracaju, SE, 49032-490, Brazil. ²Embrapa, Tabuleiros Costeiros, Aracaju, SE, 49025-040, Brazil. ³Programa de Pós-Graduação em Biotecnologia de Recursos Naturais, Universidade Federal de Sergipe, São Cristóvão, SE, 49100-000, Brazil. ⁴Embrapa, Tabuleiros Costeiros, Aracaju, SE, 49025-040, Brazil/Programa de Pós-Graduação em Biotecnologia Industrial, Universidade Tiradentes/Instituto de Tecnologia e Pesquisa, Aracaju, SE, 49032-490, Brazil.

Published: 1 October 2014

References

1. Ravikumar S, Fredimoses M, Gnanadesigan M: **Anticancer property of sediment actinomycetes against MCF-7 and MDA-MB-231 cell lines.** *Asian Pac J Trop Biomed* 2012, **2**:92-96.
2. Nawani N, Aigle B, Mandal A, Bodas M, Ghorbel S, Prakash D: **Actinomycetes: Role in Biotechnology and Medicine.** *BioMed Res Int* 2013, **2013**:1, Article ID 687190.

¹Programa de Pós-Graduação em Biotecnologia Industrial, Universidade Tiradentes/Instituto de Tecnologia e Pesquisa, Aracaju, SE, 49032-490, Brazil
Full list of author information is available at the end of the article

3. Nikolouli K, Mossialos D: **Bioactive compounds synthesized by non-ribosomal peptide synthetases and type-I polyketide synthases discovered through genome-mining and metagenomics.** *Biotechnol Lett* 2012, **34**:1393-1403.
4. Parsley LC, Linneman J, Goode AM, Becklund K, George I, Goodman RM, Lopanik NB, Liles MR: **Polyketide synthase pathways identified from a metagenomic library are derived from soil acidobacteria.** *FEMS Microbiol Ecol* 2011, **78**:176-187.
5. Ayuso-Sacido A, Genilloud O: **New PCR primers for the screening of NRPS and PKS-I systems in actinomycetes: detection and distribution of these biosynthetic gene sequences in major taxonomic groups.** *Microb Ecol* 2005, **49**:10-24.

doi:10.1186/1753-6561-8-S4-P175

Cite this article as: Hernández- Macedo *et al.*: Antimicrobial potential of *Actinomycetes* by NRPS and PKS-I pathways. *BMC Proceedings* 2014 **8** (Suppl 4):P175.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

