

### **POSTER PRESENTATION**

**Open Access** 

# Is multi-drugs resistant Acinetobacter baumannii epidemic spread related to reduced susceptibility to biocides?

B Casini<sup>1,2\*</sup>, MG Minacori<sup>1</sup>, A Buzzigoli<sup>1</sup>, P Valentini<sup>1,2</sup>, P Morici<sup>1</sup>, S Barnini<sup>3</sup>, C Tascini<sup>4</sup>, F Menichetti<sup>4</sup>, GM Rossolini<sup>5</sup>, G Privitera<sup>1,2</sup>

From International Conference on Prevention & Infection Control (ICPIC 2011) Geneva, Switzerland. 29 June – 2 July 2011

#### Introduction / objectives

Aim of this study was to assess the role of active extrusion in mediating decreased susceptibility to biocides in the persistance and spread of multi-drugs resistant of *A. baumannii* strains isolated in intensive care units of a tertiary-care teaching hospital.

#### **Methods**

67 clinical and 24 environmental strains isolated from 2007 to 2010 were genotyped by PFGE, MLST and REP-PCR. Multiplex PCRs were performed for identification of the ompA, csuE and  $bla_{\rm OXA-51like}$  sequence type groups. The antimicrobial susceptibility was determined and the presence of carbapenemase-encoding genes was analysed by characterization of the  $bla_{\rm OXA}$  genes. Chlorine susceptibility was analysed according to BS EN 1040:1997

#### **Results**

The cross-analysis of genotyping methods allowed to group strains into 4 clones, only the clone A belonging to the Group 1 corrisponding to European II clonal complex and the clone B belonging to Group 2 clonal complex European I. Since 2008, a new variant of clone A has emerged as the predominant clinical and environmental strain, resulting positive for the presence of the carbapenemase OXA-58 plasmid-mediated and showing reduced chlorine susceptibility *in vitro* 

#### Conclusion

Multi-antibiotic resistance may not be the only strategy applied by *A. baumannii* to spread and persist in health-care setting, but also the increased resistance to biocides

#### Disclosure of interest

None declared.

#### **Author details**

<sup>1</sup>Dep. Experimental Pathology, MBIE, University of Pisa, Pisa, Italy. <sup>2</sup>U.O. Igiene ed Epidemiologia, Azienda Ospedaliero-Universitaria Pisana, Pisa, Italy. <sup>3</sup>U.O. Microbiologia, Azienda Ospedaliero-Universitaria Pisana, Pisa, Italy. <sup>4</sup>U.O. Malattie Infettive, Azienda Ospedaliero-Universitaria Pisana, Pisa, Italy. <sup>5</sup>Dep. Molecular Biology, University of Siena, Siena, Italy.

Published: 29 June 2011

doi:10.1186/1753-6561-5-S6-P300

Cite this article as: Casini et al.: Is multi-drugs resistant Acinetobacter baumannii epidemic spread related to reduced susceptibility to biocides? BMC Proceedings 2011 5(Suppl 6):P300.

## Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at www.biomedcentral.com/submit



<sup>1</sup>Dep. Experimental Pathology, MBIE, University of Pisa, Pisa, Italy Full list of author information is available at the end of the article

