

POSTER PRESENTATION

Open Access

Point prevalence and risk factors of hospital acquired infections in a cluster of university affiliated hospitals in Shiraz, Iran

M Askarian^{1*}, M Yadollahi¹, O Assadian²

From International Conference on Prevention & Infection Control (ICPIC 2011)
Geneva, Switzerland. 29 June – 2 July 2011

Introduction / objectives

Hospital acquired infections (HAIs) are one of the most critical complications in hospitalized patients, responsible for a major health and economic burden. The aim of this point prevalence study of HAI was conducted in Shiraz, Iran.

Methods

The study was designed as four point prevalence surveys with identical design in eight university hospitals, each consisting of 60-700 beds, during all four seasons in 2008-2009. All patients admitted for ≥ 48 hours were studied. For all patients, a standardized data collecting form was completed, consisting of name, age, gender, presence or absence of HAI, administration of any antibiotic, insertion of central line, an endotracheal tube, mechanical ventilation, and any urinary catheter. HAI's definitions were based on the US National Nosocomial Infection Surveillance (NNIS) definitions.

Results

Data from 3450 patients were analyzed. The prevalence of HAI found to be 9.4%. The most common HAIs were blood stream infection (2.5%), surgical site infection (2.4%), urinary tract infection (1.4%), and pneumonia (1.3%). Logistic regression analysis showed that the OR of acquiring infections in males was 1.56 (95%CI 1.21-2.02), higher than in females. Other risk factors for HAI included central intravascular catheter adjusted OR 3.86 (95% CI 2.38-6.26), and urinary catheter adjusted OR 3.06 (95% CI 2.19-4.28).

Conclusion

This point prevalence study showed that HAIs are frequent in Shiraz university hospitals, and that the proportion of antibiotic prescription is high. It implies more efforts in primary prevention of HAI associated with the use of indwelling devices, and prevention of SSI.

Disclosure of interest

None declared.

Author details

¹Community Medicine, Shiraz University of Medical Sciences, Shiraz, Iran, Islamic Republic Of. ²Department for Plasma Medicine, Ernst-Moritz-Arndt University Greifswald, Walther Rathenau Strasse 49a, 17489 Greifswald, Germany.

Published: 29 June 2011

doi:10.1186/1753-6561-5-S6-P237

Cite this article as: Askarian et al.: Point prevalence and risk factors of hospital acquired infections in a cluster of university affiliated hospitals in Shiraz, Iran. *BMC Proceedings* 2011 **5**(Suppl 6):P237.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹Community Medicine, Shiraz University of Medical Sciences, Shiraz, Iran, Islamic Republic Of
Full list of author information is available at the end of the article