

## **POSTER PRESENTATION**

**Open Access** 

## Formation of woody biomass is regulated by class III HD Zip genes

Marcel Robischon

From IUFRO Tree Biotechnology Conference 2011: From Genomes to Integration and Delivery Arraial d Ajuda, Bahia, Brazil. 26 June - 2 July 2011

Secondary growth and the development of woody tissue is a key process in the formation of woody biomass. The gene family of Class III HDZip genes has been shown in the herbaceous Arabidopsis model to play a central role in regulating polarity and vascular development. While Arabidopis is a poor model for investigating processes of wood formation, in this project all poplar Class III HDZip genes were cloned and expressed in hybrid aspen as a tree model system. To circumvent an endogenous regulation mechanism involving microRNAs the sequences were also mutated to render them microRNA resistant. Lines expressing the mutated Poplar ortholog of the Arabidopsis Revoluta gene (Populus Revoluta PRE) show a spectacular phenotype with stunted growth, radialized and rolled leaves, and a double and at times triplication or quadruple layer of the xylem, suggesting the formation of multiple layers of cambium. ClassIII HDZip genes have thus been shown to be crucial for the formation of lignified tissue in trees.

Published: 13 September 2011

doi:10.1186/1753-6561-5-S7-P123

Cite this article as: Robischon: Formation of woody biomass is regulated by class III HD Zip genes. BMC Proceedings 2011 5(Suppl 7): P123

## Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at www.biomedcentral.com/submit





