

Poster presentation

Open Access

## Pro- and anti-apoptotic signaling pathways in poliovirus-infected neuronal cells

Arnaud Autret<sup>1</sup>, Sandra Martin-Latil<sup>1</sup>, Cynthia Brisac<sup>1</sup>, Laurence Mousson<sup>1</sup>, Florence Colbère-Garapin<sup>1</sup>, Jérôme Estaquier<sup>2</sup> and Bruno Blondel<sup>\*1</sup>

Address: <sup>1</sup>Biology of Enteric Viruses, Institut Pasteur, 75015 Paris, France and <sup>2</sup>INSERM V841, 94010 Créteil, France

Email: Bruno Blondel\* - [bblondel@pasteur.fr](mailto:bblondel@pasteur.fr)

\* Corresponding author

from Infectious diseases of the nervous system: pathogenesis and worldwide impact  
Paris, France. 10–13 September 2008

Published: 23 September 2008

BMC Proceedings 2008, 2(Suppl 1):P4

This abstract is available from: <http://www.biomedcentral.com/1753-6561/2/S1/P4>

© 2008 Autret et al; licensee BioMed Central Ltd.

Poliovirus (PV), the etiological agent of paralytic poliomyelitis, is a member of the genus *Enterovirus* belonging to the *Picornaviridae* family. Infection of motoneurons by PV is the direct cause of flaccid paralyzes characteristic of poliomyelitis. In PV-infected mice, motor neurons die through an apoptotic process. However, mechanisms by which PV induces cell death in neurons remain unclear. Analyses of the apoptotic pathways induced following PV infection in several cell lines have demonstrated that mitochondria are key actors of PV-induced apoptosis. We have shown that mitochondrial dysfunction in PV-infected human neuronal IMR5 cells was dependent on the protein Bax, a proapoptotic member of the Bcl-2 family. Bax activation was mediated by c-Jun NH<sub>2</sub>-terminal kinase (JNK) phosphorylation after PV infection [1]. Surprisingly, JNK activation occurred early after PV infection whereas apoptotic features were observed later in PV-infected cells. These events may involve a balance between pro- and antiapoptotic signals following PV infection. We have recently shown that, early after infection, PV also activates the phosphatidylinositol 3-kinase (PI3K)/Akt survival signaling pathway in these cells, limiting the extent of JNK activation, and thereby cell death. JNK inhibition is associated with PI3K-dependent negative regulation of the apoptosis signal-regulating kinase 1 (ASK1), which acts upstream from JNK in PV-infected IMR5 cells [2]. This survival pathway may limit the spread of PV-induced damage in the central nervous system during poliomyelitis.

### References

1. Autret A, Martin-Latil S, Mousson L, Wirotius A, Petit F, Arnoult D, Colbère-Garapin F, Estaquier J, Blondel B: **Poliovirus induces Bax-dependent cell death mediated by c-Jun NH<sub>2</sub>-terminal kinase.** *J Virol* 2007, **81**:7504-7516.
2. Autret A, Martin-Latil S, Brisac C, Mousson L, Colbère-Garapin F, Blondel B: **Early phosphatidylinositol 3-kinase/Akt pathway activation limits poliovirus-induced JNK-mediated cell death.** *J Virol* 2008, **82**:3796-3802.