

POSTER PRESENTATION

Open Access

Nephrolithiasis and nutrition in obesity

Laura Soldati^{1*}, Elena Dogliotti¹, Annalisa Terranegra^{1,2}, Tiziana Meschi³, Antonio Nouvenne³, Beatrice Prati³, Loris Borghi³, Alessandro Leone⁴, Simona Bertoli⁴, Giuseppe Vezzoli², Alberto Battezzati⁴

From Metabolism, diet and disease
Washington, DC, USA. 29-31 May 2012

Background

Obesity is a risk factor for nephrolithiasis (NL). According to an American study of a large cohort with a BMI >30, males increased the risk of NL of 30% and females of 200%. Moreover, diet plays an important role as NL risk factor, mainly in those western countries characterized by a large meat consume. This epidemiology study investigated the NL frequency in an obese Italian population, also considering the relation with metabolic syndrome and interaction with diet.

Materials and methods

Nutritional assessment included: nutritional questionnaires, anthropometry (BMI and waist circumference, body composition estimated by impedance and biochemistry (fasting glucose, serum lipids and transaminases). The presence of nephrolithiasis, osteoporosis, arterial hypertension, diabetes mellitus and metabolic syndrome were noted. Statistical analysis was performed by the SPSS software.

Results

We studied 532 obese Caucasian subjects (M/F 144/388; age 46±13.1; BMI 29.7±5.85 kg/m²). The stone formers (SF) were 41 (7.7% of the analyzed population, M 9,7%, F 6,9%; age 54.2±11.7 yrs; BMI 30.6±5.8 kg/m²). Non-stone formers (NSF) were 491 (92.3%, M/F 130/361; age 45.3±12.9 yrs [p=0.001]; BMI 29.6±5.8 kg/m²). The percentage of subjects with osteoporosis was higher in SF than in the NSF (15.79% vs 5.84%, p=0.018). In comparison with NSF, SF had higher mean waist size (101.5 vs 96.6 cm, p=0.02), systolic pressure (SF vs NSF, 130.5±15.3 vs 124.4±13.5 mmHg, p=0.007) and higher fasting serum glucose concentration (104±27.4 vs 94±15.3 mg/dL, p=0.02). Diet analysis did not show differences between SF

and NSF, except for a higher intake of butter, wine and white meat in SF.

Conclusions

Obese SF had an increased risk of osteoporosis, hypertension, diabetes and metabolic syndrome than obese NSF. This cohort of obese subjects presented a slight increase of prevalence in NL than in general Italian population that is around 5%. The prevalence of stone formers in this Italian obese population is lower than that observed in American obese subjects. This difference could be due to dissimilar genetic background or dietary habits. These findings suggest that Mediterranean diet may have a protective role against NL.

Author details

¹Dep. Medicine, Surgery, Dentistry, San Paolo Hospital, Milan, 20100, Italy. ²Nephrology and Dialysis Unit, San Raffaele Hospital, Milan, 20100 Italy. ³Dep. Clinical Sciences, University of Parma, Parma, 43121, Italy. ⁴Dep. Food Science, Technology and Microbiology, International Center for the Assessment for Nutritional Status, University of Milan, Milan, 20100, Italy.

Published: 1 June 2012

doi:10.1186/1753-6561-6-S3-P45

Cite this article as: Soldati et al.: Nephrolithiasis and nutrition in obesity. *BMC Proceedings* 2012 **6**(Suppl 3):P45.

¹Dep. Medicine, Surgery, Dentistry, San Paolo Hospital, Milan, 20100, Italy
Full list of author information is available at the end of the article