

POSTER PRESENTATION

Open Access

Ethnic differences in the association of fat and lean mass with bone mineral density in the Singapore population

AE Teo^{1,2*}, AC Ng³, K Venkataraman⁴, ES Tai⁵, YS Lee^{6,7}, EY Khoo⁵, CM Khoo⁵, SA Sadananthan^{4,6}, SS Velan^{6,8}, V Zagorodnov⁹, YS Chong⁴, P Gluckman⁶, MK Leow^{6,10}

From International Conference for Healthcare and Medical Students 2011
Dublin, Ireland. 4-5 November 2011

Introduction

Obesity and osteoporosis are two global health problems with pronounced morbidity and mortality. While body weight appears to mitigate the development of osteoporosis, whether excess body fat promotes or protects against osteoporosis remains a conundrum. The effect of ethnicity on these associations has also been understudied. We hypothesize that (1) fat mass (FM) and lean mass (LM) are independently associated with bone mineral density (BMD) and that (2) ethnic differences exist in the association of FM and LM with BMD among Chinese, Malay and Indian subjects.

Methods

We evaluated 150 overweight male subjects aged ≥ 21 years with body mass index ≥ 25 from 3 ethnic groups (Chinese =73; Malays =41; Indians =36). BMD in five regions (lumbar spine, femoral neck, total hip, ultra-distal radius and one-third radius), FM and LM were measured by dual-energy X-ray absorptiometry (DEXA) using a Hologic Discovery Wi densitometer. Whole abdomen subcutaneous and visceral fat volumes were determined by magnetic resonance imaging (MRI) and a validated segmentation algorithm. Linear regression models were developed to test the association of FM/LM with BMD, and univariate ANOVA was used to test for interaction between ethnicity and FM/LM with BMD.

Results

After adjusting for age and height, LM was positively correlated with BMD in all three ethnicities, but in different

skeletal sites: weight-bearing regions (femoral neck, hip) in Chinese, and non-weight-bearing regions (ultra-distal and one-third radius) in Malays and Indians. A negative correlation between FM and BMD was observed consistently in all regions for the Indians, especially at the hip. Visceral fat was negatively correlated with BMD, being most pronounced among Chinese and least for Malays. The interaction models revealed that with each unit of LM, Malays showed a greater increment in BMD than Chinese and Indian subjects at the ultra-distal radius.

Conclusions

Our findings suggest that FM and LM affect BMD in opposite directions, with different physiological reasons modulating this relationship. Substantial ethnic differences were observed in the association of FM and LM with BMD. These results may help explain the variation in hip and wrist fracture rates by ethnicity, and may warrant ethnic-specific clinical recommendations.

Author details

¹Agency for Science, Technology and Research, Singapore. ²University of Cambridge, UK. ³Department of Endocrinology, Singapore General Hospital, Singapore. ⁴Department of Obstetrics and Gynaecology, Yong Loo Lin School of Medicine, National University of Singapore, Singapore. ⁵Department of Medicine, Yong Loo Lin School of Medicine, National University of Singapore, Singapore. ⁶Singapore Institute for Clinical Sciences, Agency for Science, Technology and Research, Singapore. ⁷Department of Paediatrics, Yong Loo Lin School of Medicine, National University of Singapore, Singapore. ⁸Singapore Bioimaging Consortium, Agency for Science, Technology and Research, Singapore. ⁹Department of Computer Engineering, Nanyang Technological University, Singapore. ¹⁰Department of Endocrinology, Tan Tock Seng Hospital, Singapore.

¹Agency for Science, Technology and Research, Singapore
Full list of author information is available at the end of the article

Published: 9 July 2012

doi:10.1186/1753-6561-6-S4-P43

Cite this article as: Teo *et al.*: Ethnic differences in the association of fat and lean mass with bone mineral density in the Singapore population.

BMC Proceedings 2012 **6**(Suppl 4):P43.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

