

POSTER PRESENTATION

Open Access

Marine natural compounds can be efficient toward aflatoxigenic *Aspergillus flavus* strain

Yasmim Dantas Crivelenti^{1*}, Cyntia AM Arevabini¹, Mariana H De Abreu¹, Tamires Aparecida Bitencourt¹, Thaís B Mesquita¹, Bruna AM Cantelli¹, Mario FC Santos², Roberto GS Berlinck², Eduardo Hajdu³, Renê de O Beleboni¹, Mozart Marins¹, Ana Lúcia Fachin¹

From 5th Congress of the Brazilian Biotechnology Society (SBBIOTEC)
Florianópolis, Brazil. 10-14 November 2013

Background

The presence of high levels of aflatoxin is a serious problem to the production of raw peanuts and peanut crumbs. The high incidence of aflatoxin in peanuts in our country is mainly due to problems in primary production. High humidity and temperature conditions increase the likelihood of *Aspergillus* development and aflatoxins production, which is worsened during rainy weather [1]. Aflatoxins may remain in the food after the death of fungus without visible alterations [2]. The effects of aflatoxins on human and animal health, besides resistance of fungi to conventional antifungal agents has motivated the search for new inhibitors. The research of marine natural products from sponges has been considered as a promising source for the development of new antifungal agents in order to discover compounds more effective and less toxic [3]. The objective of this study was to evaluate the antifungal activity of 21 marine natural compounds toward an aflatoxigenic *A.flavus* ATCC strains.

Methods

Aflatoxin producing (CCT 7836) and non producing (ATCC9643) *A. flavus* strains were purchased from Fundação André Tosello, Campinas, Brazil and were kept in Sabouraud media at 35 °C. The minimum inhibitory concentration (MIC) of 21 sponge marine natural products, identified as SM1 to SM21, toward the two strains of *A. flavus* was determined by microdilution assay in 96-well plates using RPMI medium according to the protocol NCCLS M-38 [4] for 7 days at 37C, using commercial antifungal cercobin as control.

Results and conclusions

The marine product SM5 (CIM = 3.9 µg/mL) was more effective than cercobin (CIM = 7.8 µg/mL) against *A. flavus* control aflatoxin producer strain. Moreover, marine product SM5 (MIC = 1.9 µg/mL) was also a more effective antifungal agent than cercobin (MIC = 3.9 µg/mL) against the *A. flavus* ATCC strain. The findings suggest that natural marine products are a promising source of new molecules for the development of antifungal compounds against aflatoxigenic fungus that affect public health and the food production.

Acknowledgements

This study was supported by grants from Fundação de Amparo à Pesquisa do Estado de São Paulo, CNPq, CAPES and PET-MEC.

Authors' details

¹Unidade de Biotecnologia, Universidade de Ribeirão Preto, CEP 14096-900, Ribeirão Preto, SP, Brazil. ²Instituto de Química de São Carlos, Universidade de São Paulo, CP 780, CEP 13560-970, São Carlos, SP, Brazil. ³Museu Nacional, Universidade Federal do Rio de Janeiro, Quinta da Boa Vista, s/n, Rio de Janeiro, RJ, Brazil.

Published: 1 October 2014

References

1. Hedayati MT, Pasqualotto AC, Warn PA, Bowyer P, Denning DW: **Aspergillus flavus: human pathogen, allergen and mycotoxin producer.** *Microbiology* 2007, **153**:1677.
2. Yu J, Cleveland TE, Nierman WC, Bennett JW: **Aspergillus flavus genomics: gateway to human and animal health, food safety, and crop resistance to diseases.** *Revista Iberoamericana de Micologia* 2005, **22**:194.
3. Mayer AM, Rodriguez AD, Berlinck RG, Fusetani N: **Marine pharmacology in 2007-8: Marine compounds with antibacterial, anticoagulant, antifungal, anti-inflammatory, antimalarial, antiprotozoal, antituberculosis, and antiviral activities; affecting the immune and nervous system, and other miscellaneous mechanisms of action.** *Comparative biochemistry and physiology Toxicology & pharmacology : CBP* 2011, **153**:191-222.
4. **Reference method for broth dilution antifungal susceptibility testing of conidium-forming filamentous fungi.** *Proposed standard NCCLS document*

¹Unidade de Biotecnologia, Universidade de Ribeirão Preto, CEP 14096-900, Ribeirão Preto, SP, Brazil

Full list of author information is available at the end of the article

M38-A National Committee for Clinical Laboratory Standards, Wayne; 2002,
National Committee for Clinical Laboratory Standards.

doi:10.1186/1753-6561-8-S4-P114

Cite this article as: Criveleni *et al.*: Marine natural compounds can be efficient toward aflatoxigenic *Aspergillus flavus* strain. *BMC Proceedings* 2014 **8**(Suppl 4):P114.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

